Ultrashield[™]

SURFACE AND ENVIRONMENTAL DISINFECTANT

Composition:

Complex Formulation of Stabilized Hydrogen Peroxide I.P.
Silver Nitrate I.P.

10% w/v 0.01% w/v

Mode of Application:

For Surface Disinfection :

Surface of Hospitals, Operation Theatres, ICU's Polyclinics, Dental Clinics, OPD's Blood Banks, Clinical/ Diagnostic Laboratories.

· For Fogging:

Fogging can be done with Fogger Machine. Optimum efficacy is obtained with low pressure sprayers. A 1000 cu. ft. area room is ready in 60 minutes with only 20 % Ultrashield

Please Note:

- To reduce evaporation before & after application, close window & switch off ACs/Fans.
- Disinfected surface must remain in contact i.e. wet with solution for 20 minutes for optimum result.

Dilution Chart:

Application	Surface Disinfection	Fogging
Conc.	10% v/v	20% v/v
Ultrashield Plus	100 ml	200 ml
Potable Drinking Water	900 ml	800 ml
Contact Time	20 min.	20 min

Store protected from light and moisture in non - metallic containers





Salient Features:

- ▶ Non toxic & non irritant
- ▶ Non carcinogenic & non mutagenic
- Good material compatibility